

Agriculture and Climate Change - Adapting Crops to Increased Uncertainty (AGRI 2015)

## Selection of inbred lines and their correspondent hybrids under ultra-spaced and highly dense at normal and water-stress conditions

*Christos Dordas<sup>a\*</sup>, Fotis Gekas<sup>a</sup>, Chrysanthi Pankou<sup>a</sup>, Elissavet Ninou<sup>a</sup>, Ioannis Mylonas<sup>a</sup>, Konstantinos Tsantarmas<sup>b</sup>, Evangelia Sinapidou<sup>b</sup>, Anastasios Lithourgidis<sup>c</sup>, Iosif Sistanis<sup>d</sup>, Jovanka-Katarzyna Petrevska<sup>d</sup>, Ioannis Papadopoulos<sup>d</sup>, Paulos Zouliamis<sup>e</sup>, Anastasia Kargiotidou<sup>b</sup>, Fokion Papathanasiou<sup>d</sup>, Ioannis Tokatlidis<sup>b</sup>*

<sup>a</sup>Laboratory of Agronomy, Faculty of Agriculture, Forestry and Natural Environment, Aristotle University of Thessaloniki, 54124 Greece

<sup>b</sup>Democritus University of Thrace, Dept. of Agricultural Development, Pantazidou 193, 682 00, Orestiada, Greece

<sup>c</sup>Dept. of Agronomy, Farm of Aristotle University of Thessaloniki, 570 01 Themi, Greece

<sup>d</sup>Technological and Education Institute of Western Macedonia, Dept. of Agricultural Technology, Terma Kontopoulou, 53100 Florina, Greece

<sup>e</sup>American Genetics, Philellinon 17, 57013 Thessaloniki, Greece

### Abstract

Water stress is one of the most important environmental stresses around the world for many crop species and especially for maize. In addition, climate changes and increasing population pose serious challenges to crop improvement for increasing crop yield. Tolerance to drought is a complex quantitative trait controlled by several small effect genes or QTLs and is often confounded by differences in plants phenology. The present study aimed to investigate the tolerance of inbred lines and their correspondent hybrids using agronomic and physiological characteristics. Therefore, we used thirty one maize inbred lines under ultra-spaced and highly dense conditions and under two different water regimes (normal and drought conditions) in three different areas in Greece during the 2012 growing season. After the first year of experimentation the inbred lines were divided into three different groups tolerant, moderate sensitive and sensitive to water stress and specific hybrids from the combination of the different groups were evaluated under two water regimes and under ultra-spaced and highly dense stands and in three locations during the 2013 growing season. The inbred lines and the hybrids were evaluated using a number of physiological (gas exchange parameters, chlorophyll meter readings, chlorophyll fluorescence, water potential,

\* Corresponding author. Tel.: +30 2310 998602; fax: +30 2310 998634.

E-mail address: [chdordas@agro.auth.gr](mailto:chdordas@agro.auth.gr)

relative water content) and agronomic traits (grain yield, harvest index and yield components). The results showed that assimilation rate was reduced by the water stress at all locations and in inbred lines and also in hybrids. In addition, harvest index, instantaneous water use efficiency, and anthesis to silking interval were affected by the water stress in both the inbred lines and the hybrids. There was a clear sign that some physiological and agronomic characteristics can be used for the selection of stress-adaptive genotypes and may allow the development of new maize hybrids from specific crosses that can be grown under different conditions addressing the climate change scenarios.

© 2015 The Authors. Published by Elsevier B.V This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the organizing committee of the Agriculture and Climate Change - Adapting Crops to Increased Uncertainty (AGRI 2015)

**Keywords:** maize, drought, breeding, physiology, grain yield.

---

## References

1. Boyer JS, Byrne P, Cassman KG, et al., 2008. ‘‘Future progress in drought tolerance in maize needs new secondary traits and cross combinations’’. *The Journal of Agricultural Science* **146**:1–14.
2. Bruce WB, Edmedes GO, Barker TC, 2002. Molecular and physiological approaches to maize improvement for drought tolerance. *Journal of Experimental Botany* **53**:13–25.
3. Tokatlidis IS, 2013. Adapting maize crop to climate change. *Agronomy for Sustainable Development* **33**:63–79.

## Acknowledgements

Work co-financed by EU (ERDF) and Greek funds through the program code 09 SYN-22-604 "SYNERGASIA2009 – Action I. Cooperative small- and mid-scale projects ".